

## Original Article

## Asian Pacific Journal of Reproduction

Journal homepage: [www.apjr.net](http://www.apjr.net)

doi: 10.4103/2305–0500.284270

Effects of vascular endothelial growth factor supplementation and alginate embedding on human oocyte maturation *in vitro*Farhang Abed<sup>1</sup>, Morteza Fallah–Karkan<sup>2,3</sup>, Masoumeh Majidi Zolbin<sup>4,5</sup>, Pegah Naghizadeh<sup>1</sup>, Fereshte Aliakbari<sup>1,6</sup>, Hossein Yazdekhashti<sup>6</sup><sup>1</sup>Men's Health & Reproductive Health Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran<sup>2</sup>Laser Application in Medical Science Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran<sup>3</sup>Shohada–e–Tajrish Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran<sup>4</sup>Pediatric Urology and Regenerative Medicine Research Center, Children's Medical Center, Tehran University of Medical, Tehran, Iran<sup>5</sup>Department of Anatomy, Faculty of Medicine, Tehran University of Medical Sciences, Tehran, Iran<sup>6</sup>Department of Molecular Physiology and Biological Physics, Center for Membrane and Cell Physiology, University of Virginia, Virginia, United States

## ABSTRACT

**Objective:** To evaluate whether the use of vascular endothelial growth factor (VEGF) with alginate increases oocyte maturation following *in vitro* maturation.

**Methods:** This experimental study was performed on 150 immature oocytes (germinal vesicle oocytes) from females who were candidates for assisted reproductive technology. The germinal vesicle oocytes were randomly placed in the control, alginate, and VEGF plus alginate groups. The basic culture medium for oocytes culture (tissue culture medium 199, follicle-stimulating hormone 0.075 IU/mL, and fetal bovine serum 10%) was used in the control, alginate, and VEGF plus alginate groups. For the treatment groups (alginate, and VEGF plus alginate groups), alginate (8%) and VEGF (5 ng/mL) were added to the basic culture medium. After culture, immature oocytes were considered as oocytes unchanged in the nucleus whereas oocytes with a polar body were considered as mature oocytes (metaphase II stage). The mature oocytes in each group were fertilized by intracytoplasmic sperm injection and formed embryos were evaluated by reverse microscope.

**Results:** The oocyte maturation rate (metaphase II) significantly ( $P < 0.05$ ) increased in the alginate plus VEGF group as compared with the alginate alone and control groups during *in vitro* maturation. On day 2, the cleavage rates were significantly different in the matured oocytes between the treatment groups and the control group. The percentage of the two-cell stage, four-cell stage and eight-cell embryos was significantly higher in the treatment groups compared with the control group ( $P < 0.05$ ).

**Conclusions:** Supplementation of VEGF with alginate can improve oocyte maturation in culture media. VEGF with alginate may promote the quality of nuclear and cytoplasmic maturation of human oocytes *in vitro*.

**KEYWORDS:** Alginate; Infertility; Human oocyte; Maturation; Vascular endothelial growth factor

## 1. Introduction

The *in vitro* maturation technique is a fertility preservation method that has recently found a special place amongst the new methods of assisted reproductive technologies[1]. Favorable conditions are very important for the *in vitro* maturation technique which is used in basic biotechnology and infertility research. Oocyte maturity under laboratory conditions is a simple, hormone-free process that facilitates the study of oocyte maturation and *in vitro* fertilization[2,3]. Advantages of this technique are as follows: no need of follicle-stimulating hormone (FSH) in treatment, reduction of drug administration, and prevention of over-stimulation of the ovary in women with polycystic ovary syndrome. Also, this technique leads to transferring more embryos to compensate for lower fertility. However, the rate of pregnancy and live birth by

✉ To whom correspondence may be addressed. E-mail: [fereshtehaliakbari@yahoo.com](mailto:fereshtehaliakbari@yahoo.com); [hosein.yazdekhashti@yahoo.com](mailto:hosein.yazdekhashti@yahoo.com)

This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 4.0 License, which allows others to remix, tweak and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

**For reprints contact:** [reprints@medknow.com](mailto:reprints@medknow.com)

©2020 Asian Pacific Journal of Reproduction Produced by Wolters Kluwer- Medknow. All rights reserved.

**How to cite this article:** Abed F, Fallah-Karkan M, Majidi Zolbin M, Naghizadeh P, Aliakbari F, Yazdekhashti H. Effects of vascular endothelial growth factor supplementation and alginate embedding on human oocyte maturation *in vitro*. *Asian Pac J Reprod* 2020; 9(3): 129-134.

**Article history:** Received: 2 September 2019; Revision: 30 March 2020; Accepted: 10 April 2020; Available online: 20 May 2020

the *in vitro* maturation method using germinal vesicle (GV) oocyte obtained from the antral follicle is low[4].

The three-dimensional system, as one of the culture methods, was used for *in vitro* maturation[5]. The most important factor in a three-dimensional culture system is the selection of a suitable matrix with suitable biomechanical characteristics, high water, and gelation properties[6]. Alginate is a natural biopolymer, mainly extracted from brown algae and less frequently from bacteria. It forms 40% of the dry weight of algae and is present in the extracellular matrix of these algae, combined with calcium, magnesium and sodium cations[7]. Alginate, due to its hydrophilicity and its abundance of water properties, can provide an extracellular field material as suitable bedding for cell growth[8]. Alginate gel could support the three-dimensional architecture of the follicle. On the other hand, it can facilitate cell growth and release of food in the medium. These features make this material suitable for texture design[9].


Vascular endothelial growth factor (VEGF) is a member of the transforming growth factor  $\beta$  superfamily and it is composed of a subunit molecular in mass of 23 kDa that plays an important role in endothelial cells, ovulation and embryo implantation[10]. The most important source of VEGF in the ovarian follicle is granulosa cells[11]. Previous studies showed that adding VEGF into the *in vitro* maturation culture of bovine oocytes could promote the growth of oocyte development[12]. Some studies argued that VEGF can

increase the amount of oxygen supply in oocyte and granulosa cells and thus increase the fertilization[13,14]. Also, others believed that VEGF was directly associated with the number of injected gonadotropins and also associated with embryonic puberty[14]. Therefore, the present study was designed to investigate the effects of VEGF supplementation and alginate on human oocyte maturation *in vitro* and embryo cleavage.

## 2. Materials and methods

### 2.1. Study population

The study group consisted of women with infertility problems who referred to the fertility center of the Mahdih hospital from 2016 to 2018 in Tehran, Iran. The total number of patients consisted of 203 women with infertility problems who were under assisted reproductive technology treatment (Figure 1). Some patients underwent several *in vitro* maturation cycles during different phases of the menstrual cycle: 14 patients underwent two oocyte retrievals, 6 patients underwent three oocyte retrievals, and 1 patient underwent four oocyte retrievals. All types of infertility problems were included and maternal age was between 25 and 40 years old. The exclusion criteria were previous chemotherapy or any ovarian stimulation.



**Figure 1.** The study flowchart. VEGF: vascular endothelial growth factor.

## 2.2. Oocyte retrieval

Oocyte retrieval was done according to the center's standard for *in vitro* maturation practice 38 h after a subcutaneous administration of 10 000 IU of human chorionic gonadotropin. For retrieval of oocytes, a 19-gauge single lumen needle was used. Follicles (18–22 mm) were observed during ovulation induction. The follicular fluid including oocytes was aspirated and collected from the ovary in a 15 mL conical tube. The collected oocytes were assessed with a stereo microscope (Olympus, Japan) ( $\times 80$  magnification) and were categorized to degenerated GV, intact GV, or GV breakdown based on the presence or absence of a GV and polar body. Briefly, oocyte immaturity was observed by the presence of GV and the first polar body, and mature oocytes were assessed when the first polar body was extruded. GV oocyte with at least three layers of compact cumulus cells and with homogenous cytoplasm was selected for *in vitro* maturation. For each patient, GV oocytes were randomly placed in the control and treatment groups (alginate, and VEGF+alginate groups). Mechanical denudation of cumulus cells occurred after retrieval.

## 2.3. Preparation of alginate

Alginate solution was produced by dissolving 0.8 g of sodium alginate powder (Sigma-USA) in 100 mL of 0.9% sodium chloride solution. Then it was filtered with a 0.2  $\mu$ m filter. It was better to prepare a fresh alginate solution at any time. Alginate solution was added to 105 mM calcium chloride solution by using an insulin syringe. Alginate droplets in the calcium chloride solution were formed during 15 min, and after discharging calcium chloride, alginate seeds were added into sodium chloride (0.9%) for 10 min. After the evaporation of sodium chloride, each GV was put into alginate gel bead *via* syringe and *in vitro* maturation media were added into it.

## 2.4. In vitro maturation

Obtained GV oocytes during this study were randomly divided into three groups. The control group ( $n=50$ ) included tissue culture medium 199 (TCM 199) (Gibco), 10% fetal bovine serum (Gibco), penicillin/streptomycin 100 IU/mL (Sigma), luteinizing hormone and FSH 0.075 IU/mL (Gonal-F, Sereno); the alginate group ( $n=50$ ) included above cases with alginate 8%; the alginate+VEGF group ( $n=50$ ) included TCM 199 with alginate 8% and VEGF (5 ng/mL) (Sigma). GV oocytes in 50  $\mu$ L droplets of each group were placed in the basic culture medium and covered with mineral oil (Ovoil, Vitrolife, Sweden). Then, they were placed in a 37 °C incubator with 5% CO<sub>2</sub> for 48 h. Oocyte maturity and restoration of the meiosis were studied by a stereo microscope with  $\times 80$  magnification (Olympus, Japan) in this stage. Immature oocytes were considered as oocytes unchanged in the nucleus whereas oocytes with GV

breakdown and the absence of polar body were considered as M I oocyte. Also, oocytes with a polar body were considered as mature oocytes [metaphase (M) II stage]. In the next stage, the culture media were replaced with sodium citrate and the M II oocytes were extracted from each group for fertilization using intracytoplasmic sperm injection (her spouse's sperm).

## 2.5. Intracytoplasmic sperm injection

After the culture of oocytes for *in vitro* maturation, M II oocytes from every group were placed in 50  $\mu$ L drops that had been covered with warm mineral oil. Then, 2  $\mu$ L of the sperm were placed in a drop of polyvinyl pyrrolidone (Vitrolife, Sweden) in a plate containing a mature oocyte. And 16 to 18 h after injection of M II oocytes by processed sperm from the patient's spouse, the presence of 2 pronuclei in the oocytes fertilized was checked. Also, the number of cleavage embryos (8-cell stage) was checked on day 3 after sperm injection. The progression of embryos during *in vitro* maturation process was evaluated and none of the embryos would be transmitted inside uterus.

## 2.6. Statistical analysis

Patients' data were analyzed by analysis of variance and were expressed as mean  $\pm$  standard deviation (mean  $\pm$  SD). Statistical analyses of maturation and fertilization rates of oocytes were performed by using the *Chi*-square test in each group. Results were expressed as % and differences were considered to be significant when  $P < 0.05$ . The analyses were carried out by using SPSS Statistics for Windows, version 16 (SPSS Inc, Chicago, Ill, USA).

## 2.7. Ethical approval

The study was approved by the Ethics Committee of Shahid Beheshti University of Medical Sciences. Ethics approval number was IR.6391. All patients gave written consent for participating in the studies.

## 3. Results

### 3.1. Patients' data

There were no significant differences in age, body mass index, antral follicle counts and total oocytes between groups (Table 1).

### 3.2. Oocyte maturation

The maturity rate of GV were higher in the treat groups compared with the control group ( $P < 0.05$ ). Meanwhile, the maturity of GV was higher in the alginate+VEGF group compared with the alginate

group (Table 2; Figure 2). The rate of MII oocyte formation was higher in the alginate+VEGF group compared with the control and alginate groups.

### 3.3. Oocyte fertilization rate

On day 2, cleavage rates were significantly different between the

matured oocytes in treat groups compared with the control group ( $P<0.05$ ). The percentage of the two-cell stage and four-cell stage embryos were significantly higher in the treat groups compared with the control group ( $P<0.05$ ) (Figure 3; Table 2). The percentages of the two-cell stage, four-cell stage and eight-cell stage embryos were significantly higher in the alginate+VEGF group compared with the alginate group ( $P<0.05$ ) (Table 2).

**Table 1.** Patients' data in the control, alginate, and VEGF with alginate groups.


Parameters	Control	Alginate	Alginate+VEGF	P-value
Age (years)	32.5±1.6	31.0±3.3	32.0±3.0	0.46
Body mass index (kg/m <sup>2</sup> )	21.4±2.6	22.6±3.0	23.9±2.1	0.57
Antral follicle counts	17.5±2.0	17.5±2.0	17.5±2.3	0.49
Total oocytes	16.0±1.0	17.5±1.0	17.5±2.2	0.68

Values are expressed as mean±SD. VEGF: vascular endothelial growth factor.


**Table 2.** Comparison of maturation and fertilization rates of oocytes in the control, alginate, and VEGF with alginate groups (%).

Parameters	Control	Alginate	Alginate+VEGF	Chi value ( $\chi^2$ )	P-value
Maturation rate					
GV	16.0	11.0	9.0	27.3	0.041
M I	20.0	13.0	2.0	48.8	0.049
M II	64.0	76.0	89.0	44.9	0.039
Fertilization rate					
2-cell stage	32.0	38.5	45.0	56.8	0.049
4-cell stage	13.0	28.0	37.0	63.9	0.040
8-cell stage	-	18.0	29.0	49.9	0.048

-: No progress. VEGF: vascular endothelial growth factor; GV: germinal vesicle; M I: metaphase I; M II: metaphase II.



**Figure 2.** Maturity of human immature oocytes in metaphase I (M I) and metaphase II (M II) stages with release polar body (arrow tip) (magnification ×40). GV: germinal vesicle.



**Figure 3.** 2, 4 and 8-cell embryo from the fertilization of adult oocytes in culture medium by intracytoplasmic sperm injection (magnification ×40).

#### 4. Discussion

There are different methods for fertility preservation which can be applied based on patient's age and status as well as the risk of ovarian involvement[15,16]. The *in vitro* maturation technique is one of the fertility preservation methods[17,18].

Our study compared oocyte maturity between *in vitro* two-dimensional and three-dimensional culture systems (alginate) with VEGF involvement in. It showed that the maturity of GV oocytes is higher in the alginate system with VEGF compared with the control group.

Gonadotropin hormones were commonly used to induce oocyte maturation and ovulation in animals and humans[19]. In our study, the r-FSH hormone was used as one of the important factors in cytoplasmic and nuclear maturity of immature oocytes. Our results showed that hormones can improve the maturity of the oocytes. Similarly, Accardo *et al*[20] reported that in a culture medium containing a gonadotrophin hormone, the sheep oocyte maturation was higher.

Two-dimensional monolayer is a traditional cell culture methods. A three-dimensional cell culture is established environment for growing biological cells and are permitted to interact with their surroundings in all three dimensions. A three-dimensional cell culture permits cells *in vitro* to grow in all directions unlike two-dimensional. Three-dimensional cell culture are grown in small capsules in which the cells can grow into three-dimensional cell colonies[5]. We used two different culture systems for GV *in vitro* maturation in order to compare the efficiency of two-dimensional and three-dimensional culture methods. Although there are different two-dimensional culture systems for oocyte culturing, two-dimensional culture methods are not able to guarantee oocyte maturity and growth. Moreover, only a few studies have reported the success of oocyte maturity in two-dimensional culture methods[21]. It shows that the *in vitro* oocyte culture method is not still completely established. Different scaffolds will be evaluated for oocytes maturation applicability through three-dimensional culture. Maintaining a three-dimensional system is a crucial factor for the human clinical trial.

Alginate gel components such as fibronectin collagen I and IV have been shown to support oocyte development *in vitro*. Limited studies on the mechanical properties of alginate hydrogel and its effect on maturity oocytes have been carried out and most studies have focused on the culture of follicles in the alginate environment[22]. Xu *et al*[22] showed that by the addition of alginate to culture media, mice oocyte maturation could be successful in an alginate culture. Stouffer *et al*[23] indicated that alginate in culture media can support oocyte maturation to the MII stage. Laronda *et al*[24] suggested that human oocytes in three-dimensional coculture can provide better conditions for oocyte maturation. In the present study, the maturity of premature oocytes in alginate culture media is higher than that in the control group and our results are consistent with previous studies.

Previous studies indicated that growth factors affect oocytes maturation *in vitro* culture. In our research, we were investigated

VEGF and alginate effect on GV oocyte *in vitro*. Three-dimensional culture has been widely used recently but further studies are needed prior to clinical application. We used a 0.80% alginate system for oocyte maturation, although in previous studies 0.25% concentration of alginate was utilized for mouse follicle culture and reported that this concentration does not completely allow the entry of essential materials[25]. Matrix pore size should allow the entry of essential materials such as proteins for the maturity of the oocyte[26]. For this purpose, effective pore size should be selected; 0.80% of concentration was considered in our study.

VEGF is a growth factor involved in the formation of the embryonic circulatory system and angiogenesis. There are a few studies of the effects of VEGF on the *in vitro* maturation of human oocytes. VEGF has a variety of physiological functions in the reproductive system and folliculogenesis. It is established VEGF increased follicular vascularization and permeability of blood vessels. Also VEGF concentrations at follicular fluid increase as the antrum cavity became larger. VEGF is expressed in oocytes of follicles from all developmental stages and in the cumulus cells of antral follicles in caprine[27].

In the current study, we showed that VEGF and alginate significantly increased oocyte maturation during *in vitro* maturation of human oocytes compared with the control group. Luo *et al*[28] showed that the addition of VEGF during the *in vitro* maturation of porcine oocytes can accelerate nuclear maturation. Biswas *et al*[10] demonstrated that the addition of VEGF to porcine cumulus-oocyte complexes significantly increased embryos' developmental rate. Studies demonstrated that VEGF during *in vitro* maturation can improve the maturation oocyte, thus it may be that the beneficial effects of VEGF on oocyte *in vitro* maturation[29]. Although the maturity and growth rate of GV was significantly lower in the two-dimensional culture system. In this study, nuclear maturation was assessed by the first polar body extrusion (MII formation). Growth and maturation rates were higher in both alginate and alginate with VEGF groups[30]. We propose that VEGF might also influence oocyte maturity through the nucleus and the cytoplasm development. Further studies are required to evaluate the development of the primary embryo after intracytoplasmic sperm injection following oocyte *in vitro* maturation.

In conclusion, our results showed that *in vitro* culture conditions might influence the oocyte maturity in the presence of VEGF with alginate. VEGF with alginate can be an effective method for oocyte maturation through the nucleus and the cytoplasm development.

#### Conflict of interest statement

The authors declare that there is no conflict of interest.

#### Authors' contributions

Farhang Abed carried out the project; Morteza Fallah-Karkan and

Masoumeh Majidi Zolbin implemented the writing of the article; Pegah Naghizadeh performed the statistical analysis in this project; Fereshte Aliakbari and Hossein Yazdekhashti designed the project idea.

## References

- [1] Yang ZY, Chian RC. Development of *in vitro* maturation techniques for clinical applications. *Fertil Steril* 2017; **108**(4): 577-584.
- [2] Khalili MA, Shahedi A, Ashourzadeh S, Nottola SA, Macchiarelli G, Palmerini MG. Vitrification of human immature oocytes before and after *in vitro* maturation: A review. *J Assist Reprod Genet* 2017; **34**(11): 1413-1426.
- [3] Aliakbari F, Yazdekhashti H, Abbasi M, Hajian Monfared M, Baazm M. Advances in cryopreservation of spermatogonial stem cells and restoration of male fertility. *Microsc Res Tech* 2016; **79**(2): 122-129.
- [4] Kedem A, Yerushalmi GM, Brengauz M, Raanani H, Orvieto R, Hourvitz A, et al. Outcome of immature oocytes collection of 119 cancer patients during ovarian tissue harvesting for fertility preservation. *J Assist Reprod Genet* 2018; **35**(5): 851-856.
- [5] Haycock JW. *3D cell culture: A review of current approaches and techniques*. Vol. 695. New Jersey: Humana Press; 2011, p. 1-15.
- [6] Desai N, Abdelhafez F, Calabro A, Falcone T. Three dimensional culture of fresh and vitrified mouse pre-antral follicles in a hyaluronan-based hydrogel: A preliminary investigation of a novel biomaterial for *in vitro* follicle maturation. *Reprod Biol Endocrinol* 2012; **10**(1): 29.
- [7] Kretlow JD, Klouda L, Mikos AG. Injectable matrices and scaffolds for drug delivery in tissue engineering. *Adv Drug Deliv Rev* 2007; **59**(4-5): 263-273.
- [8] Pawar SN, Edgar KJ. Alginate derivatization: A review of chemistry, properties and applications. *Biomaterials* 2012; **33**(11): 3279-3305.
- [9] Yang JS, Xie YJ, He W. Research progress on chemical modification of alginate: A review. *Carbohydr Polym* 2011; **84**(1): 33-39.
- [10] Biswas D, Hyun SH. Supplementation with vascular endothelial growth factor during *in vitro* maturation of porcine cumulus oocyte complexes and subsequent developmental competence after *in vitro* fertilization. *Theriogenology* 2011; **76**(1): 153-160.
- [11] Cao X, Zhou P, Luo H, Zhao Y, Shi G. The effect of VEGF on the temporal-spatial change of  $\alpha$ -tubulin and cortical granules of ovine oocytes matured *in vitro*. *Anim Reprod Sci* 2009; **113**(1-4): 236-250.
- [12] Luo H, Kimura K, Aoki M, Hirako M. Effect of vascular endothelial growth factor on maturation, fertilization and developmental competence of bovine oocytes. *J Vet Med Sci* 2002; **64**(9): 803-806.
- [13] Bokal EV, Vrtovec HM, Klun IV, Verdenik I. Prolonged HCG action affects angiogenic substances and improves follicular maturation, oocyte quality and fertilization competence in patients with polycystic ovarian syndrome. *Hum Reprod* 2005; **20**(6): 1562-1568.
- [14] Van Blerkom J, Antczak M, Schrader R. The developmental potential of the human oocyte is related to the dissolved oxygen content of follicular fluid: Association with vascular endothelial growth factor levels and perifollicular blood flow characteristics. *Hum Reprod* 1997; **12**(5): 1047-1055.
- [15] Karami H, Hadad AH, Fallahkarkan M. Six years' experience of laparoscopic varicocelectomy using bipolar electrosurgery and its effect on semen parameters. *Urol J* 2016; **13**(4): 2788-2793.
- [16] Babak J, Behruz F, Mohammadreza Y, Morteza FK. The effect of human chorionic gonadotropin therapy on semen parameters and pregnancy rate after varicocelectomy. *Curr Urol* 2017; **11**(2): 92-96.
- [17] Yazdekhashti H, Rajabi Z. Established and future promising fertility preservation options in adolescence and adults. *Int Med Med Invest J* 2018; **3**(1): 1-7.
- [18] Allameh F, Abedi A, Karimi M, Fallah-Karkan M. Testicular sperm for intracytoplasmic sperm injection in oligozoospermic men with high sperm DNA fragmentation, to do or not to do. *Men's Health J* 2019; **2**(1): e4.
- [19] Wei S, Gong Z, Zhao H, Liang H, Lai L, Deng Y. Equine chorionic gonadotropin influence on sheep oocyte *in vitro* maturation, apoptosis, and follicle-stimulating hormone receptor and luteinizing hormone receptor expression. *Genet Mol Res* 2016; **15**(4): 1-12.
- [20] Accardo C, Dattena M, Pilichi S, Mara L, Chessa B, Cappai P. Effect of recombinant human FSH and LH on *in vitro* maturation of sheep oocytes; embryo development and viability. *Anim Reprod Sci* 2004; **81**(1-2): 77-86.
- [21] McLaughlin M, Patrizio P, Kayisli U, Luk J, Thomson TC, Anderson RA, et al. mTOR kinase inhibition results in oocyte loss characterized by empty follicles in human ovarian cortical strips cultured *in vitro*. *Fertil Steril* 2011; **96**(5): 1154-1159.
- [22] Xu M, West E, Shea LD, Woodruff TK. Identification of a stage-specific permissive *in vitro* culture environment for follicle growth and oocyte development. *Biol Reprod* 2006; **75**(6): 916-923.
- [23] Xu J, Lawson M, Yeoman R, Pau K, Barrett S, Zelinski M, et al. Secondary follicle growth and oocyte maturation during encapsulated three-dimensional culture in rhesus monkeys: effects of gonadotrophins, oxygen and fetuin. *Hum Reprod* 2011; **26**(5): 1061-1072.
- [24] Laronda MM, Duncan FE, Hornick JE, Xu M, Pahnke JE, Whelan KA, et al. Alginate encapsulation supports the growth and differentiation of human primordial follicles within ovarian cortical tissue. *J Assist Reprod Genet* 2014; **31**(8): 1013-1028.
- [25] Shikanov A, Xu M, Woodruff TK, Shea LD. Interpenetrating fibrin-alginate matrices for *in vitro* ovarian follicle development. *Biomaterials* 2009; **30**(29): 5476-5485.
- [26] West ER, Xu M, Woodruff TK, Shea LD. Physical properties of alginate hydrogels and their effects on *in vitro* follicle development. *Biomaterials* 2007; **28**(30): 4439-4448.
- [27] Araújo VR, da Silva GM, Figueiredo JR. Only VEGF is necessary to stimulate the meiotic competence of caprine oocytes during *in vitro* maturation. *Ciência Anim* 2017; **27**(1): 31-40.
- [28] Luo H, Kimura K, Aoki M, Hirako M. Vascular endothelial growth factor (VEGF) promotes the early development of bovine embryo in the presence of cumulus cells. *J Vet Med Sci* 2002; **64**(11): 967-971.
- [29] Anchordoquy JM, Anchordoquy JP, Testa JA, Sirini MÁ, Furnus CC. Influence of vascular endothelial growth factor and cysteamine on *in vitro* bovine oocyte maturation and subsequent embryo development. *Cell Biol Int* 2015; **39**(10): 1090-1098.
- [30] Higuchi CM, Maeda Y, Horiuchi T, Yamazaki Y. A simplified method for three-dimensional (3-D) ovarian tissue culture yielding oocytes competent to produce full-term offspring in mice. *PLoS One* 2015; **10**(11): e0143114.