



# Prevalence of Sexually Transmitted Diseases in Men of 25 - 60 Years in Iran

Fereshteh Aliakbari<sup>1</sup>, Majid Emadeddin<sup>1</sup>, Neda Taghizabet<sup>1</sup>, Fatemeh Rezaei-Tazangi<sup>2</sup>, Zahra Sadeghzadeh<sup>1</sup> and Jalil Hosseini<sup>1,\*</sup>

<sup>1</sup>Men's Health and Reproductive Health Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

<sup>2</sup>Department of Anatomical Sciences, Faculty of Medicine, Fasa University of Medical Sciences, Fasa, Iran

\*Corresponding author: Men's Health and Reproductive Health Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran. Email: senmohr1368@gmail.com

Received 2021 July 10; Accepted 2021 July 29.

## Abstract

**Background:** Sexually transmitted diseases (STDs), especially in men, can lead to multiple morbidities and facilitate the transmission of serious pathogens, such as the human immunodeficiency virus. The present study used the data from a nationwide survey on male morbidities to estimate the prevalence of STDs among men in Iran.

**Methods:** This cross-sectional study used the data from a nationwide project on male reproductive morbidities in 2007. A total of 2296 men within the age range of 25 - 60 years were recruited by systematic cluster sampling from four provinces (i.e., Golestan, Hormozgan, Kermanshah, and Isfahan, Iran). The data on the symptoms of STDs, including genital secretion, pushing out, itching or genital ulcers, and lymphadenopathy of the inguinal area after sexual contact, were collected by trained urologists. Data analysis was carried out using SPSS software (version 18.0, Chicago, IL, USA). The independent samples *t*-test and chi-square test were used for data analysis.

**Results:** A total of 2296 men with a mean age value of  $39.95 \pm 10.3$  years were interviewed in this study. Two-thirds of all the subjects (75%) were aware of using condoms in suspected sexual relationships; however, only 69% of the aforementioned subjects used condoms in those circumstances. Overall, 14 subjects (0.6%) had one type of STDs at the study time. Moreover, 110 subjects (4.7%) answered "yes" to the question of whether they were referred to a physician for sexual problems.

**Conclusions:** The prevalence of STDs is still low among the male population in Iran. Careful and well-designed surveillance systems to monitor the incidence of STDs and proper preventive measures to restrict the spread of sexually transmitted pathogens are among the proposed recommendations.

**Keywords:** Sexually Transmitted Diseases, Male, Iran, Population-Based Survey

## 1. Background

Sexually transmitted diseases (STDs), such as gonorrhea, chlamydia infection, syphilis, genital herpes, and the human immunodeficiency virus/acquired immunodeficiency syndrome, are common and serious health problems worldwide (1). The STDs impose a heavy burden on health care systems and are attributed to serious health conditions in both genders. The importance of STDs is such that the World Health Organization lists *Neisseria gonorrhoeae* as one of twelve organisms on the "global priority list of antibiotic-resistant bacteria" (2).

Approximately 126 million new cases of STDs annually occur; nevertheless, two-thirds of STDs cases are reported or diagnosed among adolescents and young adults (3). Global statistics show that annually one-third of new cases of STDs are under 25 years of age (4). The STDs also account

for most male urethritis; nonetheless, the symptoms include urethral discharge, irritation, and dysuria (5). In men younger than 35 years of age, *Chlamydia trachomatis* is the most common cause of epididymitis (6). Furthermore, comorbidities are common among STD pathogens. For instance, epididymitis, which causes pain and swelling almost always unilateral, can be accompanied by inflammation in the testes, especially viral orchitis (7, 8).

Despite the magnitude and seriousness, the implementation of primary and secondary prevention against STDs is not straightforward in many settings. For example, the primary prevention of STDs is typically orchestrated by training and educating on the most important mode of transmission (i.e., sexual relationship) (9). Nevertheless, educating on safe sex is not socially acceptable occasionally and might arise various social and even political

confrontations. Despite this obstacle, STDs can be potentially controlled by several well-established activities, such as partner notification, which involves interviewing individuals diagnosed with STDs, obtaining information about their sexual partners, and locating these partners so that they can be referred for testing and treatment (10, 11).

Regarding secondary prevention, the emergence of antimicrobial resistance and inherent behaviors of some STD pathogens to cause asymptomatic infection, especially in men, might obstruct the early detection and timely treatment of STDs. A survey conducted on 16 sexual health clinics in England reported a gender difference in treatment-seeking behavior following the contraction of asymptomatic STDs; accordingly, seeking treatment by asymptotically infected men with STDs is expected only when there was a symptomatic disease or diagnosis in their partners (12).

Undoubtedly, sexual health has great importance on the human quality of life; however, scientific evidence on sexual health and morbidities from developing countries is scarce. On the other hand, the prioritization of reproductive health problems and resource allocation to address priority issues require valid epidemiological information. For the provision of such data, a comprehensive nationwide project was carried out to identify the prevalence and contributory factors of three reproductive morbidities (namely infertility, STDs, and sexual dysfunction) in the male population within the age range of 25 - 60 years in Iran. The present study used a part of the aforementioned data to estimate the prevalence of STDs and determine important related factors to this problem.

## 2. Methods

This cross-sectional study used the data from a nationwide project entitled "National Survey On Reproductive Morbidities Among Males in Iran in 2007", conducted by the Men's Health and Reproductive Health Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran, in cooperation with the United Nations Population Fund and the Iranian Ministry of Health. The study population consisted of all men within the age range of 25 - 60 years residing in four provinces across the country, including Golestan in the north, Hormozgan in the south, Kermanshah in the west, and Isfahan in the center of Iran. The selected provinces were randomly chosen from 31 provinces in Iran; however, three cities were selected from each province, and the sample size was assigned proportionally to the census population of 25 - 60-year-old males stratified on residence area (i.e., urban or rural).

Subject recruitment was carried out by systematic cluster sampling. For reaching the final sample size of 2296, 328 cluster heads (each with a size of 7) were defined based

on the registered household number in healthcare centers. After the selection of households within each cluster, one male subject from each household within the age range of 25 - 60 years was invited to visit the healthcare center. The inclusion criteria were defined as the age range of 25 - 60 years and residence in the study area for at least 3 months before conducting the study. The exclusion criteria were defined as hospitalization during the study course and unwillingness to participate in the study.

The data were collected using structured questionnaires by trained interviewers who were local general practitioners. Clinical examination included the measurement of height, weight, and blood pressure. Laboratory findings included semen analysis, urethral smear, follicle-stimulating hormone, and testosterone level. Ethical approval was granted by the Review Board of Shahid Beheshti University of Medical Sciences. Verbal informed consent was sought from the respondents before the interview.

A structured checklist, including demographic information, was filled out by the subjects. Furthermore, the data on the symptoms of STDs, including genital secretion, pushing out, itching or genital ulcers, and lymphadenopathy of the inguinal area after sexual contact, were collected by trained urologists. Interviews were performed by trained interviewers who received one training session at the Men's Health and Reproductive Health Research Center in February 2007 in Tehran. Data analysis was carried out using SPSS software (version 18.0, Chicago, IL, USA). The independent samples t-test and Chi-square test were used for data analysis.

## 3. Results

In total, 2296 men with a mean age value of  $39.95 \pm 10.3$  years were interviewed in this study. Two-thirds of the subjects (75.5%) resided in urban areas. Most subjects (90.8%) were married; however, the mean age value at first marriage was  $24.22 \pm 3.94$  years (range: 16 - 54 years). Moreover, the mean length value of marriage was  $16.92 \pm 10.5$  years. Among those who reported being single ( $n = 210$ ), 42 subjects (20%) reported having sexual intercourse, mainly with a fixed partner. Extramarital relationship, mainly with a fixed partner, was reported by 81 married subjects (4%). Two-thirds of all the subjects (75%) were aware of using condoms in suspected sexual relationships; nonetheless, only 69% of the aforementioned subjects used condoms in those circumstances.

Most subjects (80%) were reported as employed; nevertheless, the highest (9.2%) and lowest (3.6%) unemployment rates were observed in Kermanshah and Isfahan provinces, respectively. Regarding cigarette smoking, 7 and 25% of the subjects smoked occasionally and daily over

the preceding year, respectively. Daily and occasional exposures to cigarette smoking were reported by 9 and 18.5% of the subjects, respectively. Additionally, 10.4% of the subjects reported any drug abuse in occasional (7%) and daily (4%) forms. No history of STDs, including secretion from the duct, burning duct, and testicular swelling and pain, was reported by 1.4, 4.9, and 1.6% of the subjects, respectively.

### 3.1. STD Referral and Treatment

Based on the medical decision by the study interviewers as certified general practitioners, a total of 14 subjects (0.6%) had one type of STDs at the study time. Furthermore, 110 subjects (4.7%) answered "yes" to the question of whether they were referred to a physician for sexual problems, only 27% (n=30) of whom sought medical treatment. Delayed referral, defined as seeking medical treatment at least 2 weeks after the appearance of symptoms, was reported by 56.7% of symptomatic cases (n = 17).

The prevalence of STDs was higher among unmarried (1%) and employed (0.8%) subjects. The differences in the prevalence of STDs according to marital and employment status were not statistically significant. On the other hand, a significantly higher proportion (0.8%) of urban respondents reported having STDs, compared to that (0.2%) of rural respondents ( $P = 0.032$ ). Moreover, a significantly higher proportion (17.8%) of respondents with multiple sexual partners reported having STDs, compared to that (0%) of respondents with a single sexual partner ( $P = 0.012$ ). Nearly 1% of the subjects with enough knowledge of using condoms and 1% of subjects who were condom users during suspected sexual intercourse were observed to have STDs, which was significantly higher, compared to that (0%) of subjects with insufficient knowledge of the issue ( $P = 0.033$ ; Table 1).

## 4. Discussion

STDs are considered health problems in most parts of the world. The so-called term of STDs refers to a variety of clinical syndromes caused by pathogens that can be acquired and transmitted mainly through sexual activities. STDs are among the first 10 causes of unpleased diseases in young male adults. It is estimated that each year 340 million cases of STDs occur in men and women aged 15 - 49 years worldwide. More than 1 million sexually transmitted infections are acquired every day worldwide (13). The STDs have been reported from all over the world, and Iran is no exception in this regard; however, 61% of the Iranian population are in a sexually active life period (age range: 15 - 49 years) (14).

The results of the present study showed that 0.8% of employed participants had STDs; nonetheless, none of the unemployed subjects was observed to have STDs; however, the difference was not statistically significant. Regarding the relationship between occupation and STDs, Alema et al., in a study conducted on 628 individuals aged 25 - 64 years suggested that unemployment is an underlying factor for high-risk sexual behaviors, which might lead to the contraction of STDs (15). The present study showed that there was no relationship between STDs and alcohol consumption; accordingly, an equal proportion of subjects, stratified based on alcohol consumption, were observed to have STDs (0.7%).

Another well-known risk factor for STDs contraction is having multiple sexual partners. In the current study, a significantly higher proportion (18%) of subjects with multiple sexual partners reported having STDs, compared to that (0%) of subjects with a single sexual partner (0%). Several reports across the world have also indicated a relationship between the use of condoms and the risk of STDs. Koss et al. conducted a systematic review on 12 studies focusing on condom use and syphilis risk, indicating a lower probability of syphilis contraction following condom use in sexual intercourse (16). Nevertheless, further studies are needed to determine the association between condom use and syphilis prevalence.

The present study showed that the prevalence of STDs was significantly higher in individuals who used condoms in suspected sexual intercourse and had enough knowledge of condom use in suspected sexual intercourse. This finding is in contrast to the finding of previous reports showing that the consistent and correct use of male condoms is the primary method preventing the acquisition or transmission of STDs (17-19). This inconsistency might be due to the fact that condom use was reported only by subjects who reported having multiple sexual partners. In other words, condom use during sexual intercourse was not reported by any of the respondents who had a single sexual partner. This result is consistent with the results of a qualitative study reporting that condom use by men greatly depends on relationship duration. In other words, men feel that their role should be to support and communicate with their partners in a stable relationship with a single sexual partner. Therefore, an unlikely use of condoms is frequently reported in this regard. On the other hand, occasional sexual intercourse with multiple partners impose the feeling of self-protection, which leads to more likelihood of condom use by men (20).

Effective interventional programs to eliminate STDs require systematic monitoring of disease incidence and high-risk behavior prevalence, especially in young individuals. This monitoring should be accompanied by the con-

**Table 1.** Prevalence of Sexually Transmitted Diseases According to General Characteristics of Men Aged 25 - 60 Years in Iran

Variables	No. (%)	STDs		P-Value
		Absent; No. (%)	Present; No. (%)	
<b>Current marital status</b>				0.403
Married	1861 (90.3)	1849 (99.3)	12 (0.7)	
Unmarried	201 (9.7)	199 (99)	2 (1)	
<b>Employment status</b>				0.206
Employed	1850 (89.8)	1836 (99.2)	14 (0.8)	
Unemployed	210 (10.2)	210 (100)	0 (0)	
<b>Residence area</b>				0.032
Urban	1725 (75.5)	1712 (99.2)	13 (0.8)	
Rural	558 (24.5)	557 (99.8)	1 (0.2)	
<b>Diabetes status</b>				0.627
Healthy	1924 (93.2)	1911 (99.3)	13 (0.7)	
Diabetic	140 (6.8)	139 (99.2)	1 (0.8)	
<b>Drug abuse</b>				0.570
No	1852 (89.6)	1839 (99.3)	13 (0.7)	
Yes	213 (10.4)	211 (99.5)	1 (0.5)	
<b>Alcohol consumption</b>				0.633
No	2133 (93.7)	1909 (99.3)	13 (0.7)	
Yes	142 (6.3)	141 (99.3)	1 (0.7)	
<b>Type of sexual partner<sup>a</sup></b>				0.012
Multiple	45 (60)	37 (82.2)	8 (17.8)	
Fixed	30 (40)	30 (100)	0 (0)	
<b>Knowledge of using condoms</b>				0.033
Absent	455 (21.9)	455 (100)	0 (0)	
Present	1620 (78.1)	1606 (99.1)	14 (0.9)	
<b>Condom use in suspected sexual intercourse</b>				0.021
No	517 (25.5)	517 (100)	0 (0)	
Yes	1510 (74.5)	1497 (99.1)	13 (0.9)	

Abbreviation: STDs, sexually transmitted diseases.

<sup>a</sup> Among those with reported extramarital sexual relationships.

sistent application of well-designed and reliable epidemiologic methods, metrics, and progress indicators. Additionally, it is required to design accurate, precise, and easily understood monitoring tools. Based on this perspective, the implementation of primary prevention programs to educate the population, especially young individuals, on protection against STDs and the provision of safety measurements (e.g., mechanical contraception devices) are highly warranted. Based on an epidemiologic perspective, the surveillance of STDs should be strengthened to detect inherent biases, allow for fast-tracking schemes, and act and respond timely and properly. One desired attribute of such a surveillance system would be using a combination of relative and absolute measures, such as rate ratios, rate differences, and population-attributable proportions instead of any single measure in assessing the progress of prevention programs (21).

This study demonstrated that overall 0.6% of the subjects (n = 14) had at least one type of STDs at the study time.

The estimated prevalence of STDs was mainly reported among specific subpopulations, such as homosexual men, homeless individuals, and individuals with high-risk sexual activities. To the best of our knowledge, a limited number of studies have been performed to estimate the prevalence of STDs in the general population, which is one of the strengths of the present study. On the other hand, the measurement of STDs was assessed by general physicians, where some detailed examination to detect STDs might be missed. Furthermore, due to the executive inability to use laboratory diagnostic kits, it was not possible to detect the co-infections of sexually transmitted pathogens in respondents.

#### 4.1. Conclusion

The prevalence of STDs is still low among the male population in Iran. Careful and well-designed surveillance systems to monitor the incidence of STDs and proper preventive measures to restrict the spread of sexually transmitted

pathogens are among the proposed recommendations.

## Footnotes

**Authors' Contribution:** MA and JH, project development; ME, data collection; FA, SN, NT, and FRT, manuscript writing. All the authors read and signed the final manuscript.

**Conflict of Interests:** The authors declare that there is no conflict of interest.

**Ethical Approval:** Ethical approval was granted by the Review Board of Shahid Beheshti University of Medical Sciences.

**Funding/Support:** None to declare.

**Informed Consent:** All the participants entered the study with their consent.

## References

1. Workowski KA, Bolan GA; Centers for Disease Control Prevention. Sexually transmitted diseases treatment guidelines, 2015. *MMWR Recomm Rep*. 2015;**64**(RR-03):1-137. [PubMed: 26042815]. [PubMed Central: PMC5885289].
2. Shrivastava SRB, Shrivastava PS, Ramasamy J. World health organization releases global priority list of antibiotic-resistant bacteria to guide research, discovery, and development of new antibiotics. *J Med Soc*. 2018;**32**(1):76. doi: 10.4103/jms.jms\_25\_17.
3. Widman L, Nesi J, Kamke K, Choukas-Bradley S, Stewart JL. Technology-based interventions to reduce sexually transmitted infections and unintended pregnancy among youth. *J Adolesc Health*. 2018;**62**(6):651-60. doi: 10.1016/j.jadohealth.2018.02.007. [PubMed: 29784112]. [PubMed Central: PMC5966833].
4. Fineberg HV. The social dimensions of AIDS. *Sci Am*. 1988;**259**(4):128-34. doi:10.1038/scientificamerican1088-128. [PubMed: 3072669].
5. Greer SL. The European Centre for Disease Prevention and Control: Hhub or hollow core? *J Health Polit Policy Law*. 2012;**37**(6):1001-30. doi: 10.1215/03616878-1813817. [PubMed: 22899831].
6. Choudhri Y, Miller J, Sandhu J, Leon A, Aho J. Chlamydia in Canada, 2010-2015. *Can Commun Dis Rep*. 2018;**44**(2):49-54. doi: 10.14745/ccdr.v44i02a03. [PubMed: 29770099]. [PubMed Central: PMC5864312].
7. Patel MR, Brooks JT, Tie Y, Garg S, Bradley H. Prevalence of gonorrhea and chlamydia testing by anatomical site among men who have sex with men in HIV medical care, United States, 2013-2014. *Sex Transm Dis*. 2018;**45**(1):25-7. doi: 10.1097/OLQ.0000000000000691. [PubMed: 28876291]. [PubMed Central: PMC6945295].
8. Ryan L, Daly P, Cullen I, Doyle M. Epididymo-orchitis caused by enteric organisms in men > 35 years old: beyond fluoroquinolones. *Eur J Clin Microbiol Infect Dis*. 2018;**37**(6):1001-8. doi: 10.1007/s10096-018-3212-z. [PubMed: 29450767].
9. Mirzazadeh A, Biggs MA, Viitanen A, Horvath H, Wang LY, Dunville R, et al. Do school-based programs prevent HIV and other sexually transmitted infections in adolescents? A systematic review and meta-analysis. *Prev Sci*. 2018;**19**(4):490-506. doi: 10.1007/s11121-017-0830-0. [PubMed: 28786046].
10. Van den Broucke S. Strengthening public health capacity through a health promotion lens. *Health Promot Int*. 2017;**32**(5):763-7. doi: 10.1093/heapro/dax064. [PubMed: 28992188].
11. Chesson HW, Ludovic JA, Berruti AA, Gift TL. Methods for sexually transmitted disease prevention programs to estimate the health and medical cost impact of changes in their budget. *Sex Transm Dis*. 2018;**45**(1):2-7. doi: 10.1097/OLQ.0000000000000747. [PubMed: 29240632]. [PubMed Central: PMC5905677].
12. Aicken CRH, Wayal S, Blomquist PB, Fabiane SM, Gerressu M, Hughes G, et al. Pathways to, and use of, sexual healthcare among Black Caribbean sexual health clinic attendees in England: evidence from cross-sectional bio-behavioural surveys. *BMC Health Serv Res*. 2019;**19**(1):668. doi: 10.1186/s12913-019-4396-3. [PubMed: 31533716]. [PubMed Central: PMC6749649].
13. Holcombe E, Ryan S, Manlove J. Discussions about contraception or STDs prior to sex. Fact sheet. Publication# 2008-14. *Child Trends*. 2008.
14. Shaditalab J. Iranian women: Rising expectations. *Critiq Crit Middle E Stud*. 2005;**14**(1):35-55. doi: 10.1080/10669920500057005.
15. Alemu H, Mariam DH, Belay KA, Davey G. Factors predisposing out-of-school youths to HIV/AIDS-related risky sexual behaviour in northwest Ethiopia. *J Health Popul Nutr*. 2007;**25**(3):344-50. [PubMed: 18330068]. [PubMed Central: PMC2754028].
16. Koss CA, Dunne EF, Warner L. A systematic review of epidemiologic studies assessing condom use and risk of syphilis. *Sex Transm Dis*. 2009;**36**(7):401-5. doi: 10.1097/OLQ.0b013e3181a396eb. [PubMed: 19455075].
17. Eaton DK, Kann L, Kinchen S, Shanklin S, Ross J, Hawkins J, et al. Youth risk behavior surveillance - United States, 2009. *MMWR Surveill Summ*. 2010;**59**(5):1-142. [PubMed: 20520591].
18. Sanders SA, Reece M, Herbenick D, Schick V, Dodge B, Fortenberry JD. Condom use during most recent vaginal intercourse event among a probability sample of adults in the United States. *J Sex Med*. 2010;**7** Suppl 5:362-73. doi: 10.1111/j.1743-6109.2010.02011.x. [PubMed: 21029391].
19. Yarber WL, Crosby RA, Graham CA, Sanders SA, Arno J, Hartzell RM, et al. Correlates of putting condoms on after sex has begun and of removing them before sex ends: A study of men attending an urban public STD clinic. *Am J Mens Health*. 2007;**1**(3):190-6. doi: 10.1177/1557988307301276. [PubMed: 19482797].
20. Van der Heijde CM, Veldhoven M, Vonk P. How do male students perceive their role in contraception and the protection against STDs? *Eur J Public Health*. 2011;**29**(Suppl 4):165-86.
21. Dean HD, Truman BI, Mermin JH. Monitoring progress in reducing disparities in HIV, hepatitis, STDs, and tuberculosis. *Am J Public Health*. 2018;**108**(S4). S236. doi: 10.2105/AJPH.2018.304811. [PubMed: 30383420]. [PubMed Central: PMC6215374].